

Clinical Pharmacology Considerations for Evaluating Race, Ethnicity, Geography, and Ancestry During Drug Development and Regulatory Review

Anuradha (Anu) Ramamoorthy, PhD Master Scientist/Policy Lead, GPT, OCP

FDA-ISoP Workshop

Oct 16, 2023

Disclaimer: This presentation reflects the views of the speaker and should not be construed to represent FDA's views and policies

Topics for Discussion



Some Definitions: Race & Ethnicity



Self-reported and are considered social-political constructs that should not be interpreted as being scientific or anthropological in nature Ethnicity

- Hispanic or Latino
- Not Hispanic or Latino

Race

- American Indian or Alaska Native
- Asian
- Black or African American
- Native Hawaiian or Other Pacific Islander
- White

Ethnicity: Social group with a shared history, heritage, culture, sense of identity, and territorial identity that occurs despite racial dissimilarity

Hispanic or Latino: A person of Cuban, Mexican, Puerto Rican, South or Central American, or other Spanish culture or origin, regardless of race. The term, "Spanish origin," can be used in addition to "Hispanic or Latino."

Race: Groups based on physical traits regarded as common among people of shared ancestry

American Indian or Alaska Native: A person having origins in any of the original peoples of North and South America (including Central America), and who maintains tribal affiliation or community attachment.

Asian: A person having origins in any of the original peoples of the Far East, Southeast Asia, or the Indian subcontinent, including, for example, Cambodia, China, India, Japan, Korea, Malaysia, Pakistan, the Philippine Islands, Thailand, and Vietnam.

Black or African American: A person having origins in any of the black racial groups of Africa. Terms such as "Haitian" or "Negro" can be used in addition to "Black or African American."

Native Hawaiian or Other Pacific Islander: A person having origins in any of the original peoples of Hawaii, Guam, Samoa, or other Pacific Islands. White: A person having origins in any of the original peoples of Europe, the Middle East, or North Africa.

Race and ethnicity data collection guidance ; Image source: cancer.gov

Some Definitions: Geography



Some Definitions: Ancestry



Ancestry: Biologic or genetic ancestry of an individual and in genetic research studies, ancestry information is replacing the use of racial categories. Note: the term is subject to a variety of interpretations and can be based on geographic, historical, cultural, or religious definitions

Image source: https://www.alamy.com/stock-photo/, https://www.statnews.com/2022/05/02/substituting-genetic-ancestry-for-race-in-research-not-so-fast/; Jorde LB & Bamshad MJ, 2020

Some Definitions: Intrinsic & Extrinsic Factors



Intrinsic & Extrinsic Factors: Factors that can lead to exposure or response difference across subpopulations or impact the interpretation of R/B across subpopulations

Case Example: Multiple Myeloma



Analysis of racial and ethnic disparities in multiple myeloma US FDA drug approval trials

blood advances 22 MARCH 2022 - VOLUME 6, NUMBER 6

Bindu Kanapuru,^{1,*} Laura L. Fernandes,^{2,*,†} Lola A. Fashoyin-Aje,³ Andrea C. Baines,¹ Vishal Bhatnagar,⁴ Rachel Ershler,¹ Thomas Gwise,² Paul Kluetz,⁴ Richard Pazdur,⁴ Elizabeth Pulte,⁵ Yuan-Li Shen,⁶ and Nicole Gormley¹

Cancer. 2020 January 01; 127(1): 82-92. doi:10.1002/cncr.33208.

African Americans with translocation t(11;14) have superior survival after autologous hematopoietic cell transplantation for multiple myeloma compared in Whites in the United States

Talha Badar, MD¹, Parameswaran Hari, MD, MS², Omar Dávila, MPH², Raphael Fraser,

652.MULTIPLE MYELOMA AND PLASMA CELL DYSCRASIAS: CLINICAL AND EPIDEMIOLOGICAL | NOVEMBER 15, 2022

Current Testing Practices for t(11;14) Rearrangements in Patients with Newly Diagnosed Multiple Myeloma in the United States

Linda B. Baughn, Elizabeth S. Mearns, Michael Nixon, Vivek S. Chopra, Allicia Girvan, R. Frank Cornell, Jayeta Saxena, Bethany M. Slifko, Jordan Clark, Shaji K Kumar

Case Example: EGFR TKIs





Sharma VS et al., 2007; Calvayrac O et al., 2017; Ramamoorthy A et al., 2018

Differences Reported for FDA-approved New Drugs





428 new drugs approved between 2008 and 2019; ¹>20% change in AUC, Cmax, etc.; ² Genotype/phenotype frequency differences across races and ethnicities (e.g., CYP2D6, CYP2C19, NAT2, UGT1A1); ³ specific dosage or usage (monitoring, contraindication) recommendations

Modified from Ramamoorthy A et al., 2015 and Ramamoorthy A et al., 2022

30

Dose Differences Across Regions





Dose differences potentially driven by PK/PD differences, safety concerns, and other factors (e.g., business decisions)

Causes for Differences Among Subpopulations FDA



Ramamoorthy A et al., 2015





Objective

To evaluate generalizability of risk/benefit data available from the drug development program to a diverse US **population** that is expected to receive the drug upon its approval

Based on our understanding of the disease, drug, and target is a clinically relevant difference expected in drug exposure and/or response among clinically relevant subpopulations?



report subgroup data as outlined in FDA guidances and regulations¹³



Differences Among Subpopulation Can Lead to Differences in Regulatory Recommendations



Altered benefit/risk

Geographic differences in drugs or doses approved/withdrawn

Targeted recommendations across subpopulations

Subpopulations Differences Can Lead to Differences in Regulatory Recommendations



No dose adjustment

No specific monitoring recommendations



Different dosing recommendations -

Considerations for therapy selection

Warnings and Precautions

 \longrightarrow

Rosuvastatin in Asians

• Tacrolimus in Blacks

 Eliglustat based on CYP2D6 metabolizer status

ACE inhibitors (e.g., losartan, cadesartan) in Blacks

Carbamazepine for HLAB*1502 in Asians



Indication Contraindication BiDil for self-identified Blacks

 Rasburicase for G6PD deficiency

Postmarketing Studies to Obtain Additional Information on Safety, Efficacy, or Optimal Use



Conduct a randomized, controlled clinical trial to evaluate the efficacy and safety in **African-American patients** with systemic lupus erythematosus (belimumab, 2011) Evaluate the safety and PK in patients with wild-type, heterozygous, and homozygous **UGT1A1*28 genotypes** (belinostat, 2014)

Conduct an epidemiologic study to evaluate the incidence of angioedema in **Black patients** (sacubitril and valsartan, 2015) Include an assessment of PK, PD, safety, and efficacy in U.S. racial and ethnic minority patients including Black and Asian patients (daratumumab and hyaluronidase, 2020)

Summary



Diversity is often limited in clinical trials **Disease burden may be high** for some disease in some populations

Need for **continued effort to enroll underrepresented populations** during drug development

Understand exposure and/or response across clinically relevant subpopulations

Leverage clinical pharmacology to understand and evaluate the impact of intrinsic and extrinsic factors on drug exposure and/or response

