



[Student Name, DOB, Student ID]

NAYZILAM Nasal Spray Instructions

For Use Only By Trained Staff

When to Use: *[EDIT FOR INDIVIDUAL ORDERS]* Administer NAYZILAM 5 mg, using 1 nasal spray device, if a convulsive seizure lasts longer than 5 minutes or if the student experiences a cluster of 3 or more seizures within 60 minutes, per provider instructions.

Medication Location: *[EDIT LOCATION FOR THIS STUDENT]* e.g., health room, classroom, student backpack if authorized for self-carry, etc).

Confirm the 6 Rights of Medication Administration: *[EDIT FOR INDIVIDUAL ORDERS]*

Right Student	Student Name
Right Medication	NAYZILAM (midazolam) nasal spray
Right Dose	5 mg dose delivered from 1 device <i>The 5 mg and 10 mg doses are given using one device. The 15 mg and 20 mg doses are given using two devices—one device in each nostril.</i>
Right Route	Intranasal
Right Time	Per health care provider prescription
Right Documentation	Document medication administration promptly and accurately on Medication Administration Record

Equipment: Prefilled medication device(s) containing NAYZILAM nasal spray

PROCEDURE

Step 1: Peel open the blister packaging

- Do not open the blister packaging until ready to use.
- Hold the blister packaging in the palm of your hand, find the “Peel Here” tab on the foil backing, and carefully remove the device.

Step 2: Hold the nasal spray unit

- Hold the nasal spray device with your thumb on the plunger and your middle and index fingers on either side of the nozzle.
- **Do not press the plunger yet. If you press the plunger now, you will lose the dose.**

Step 3: Place the tip into 1 nostril

- Place the tip of the nozzle into 1 nostril until your fingers on either side of the nozzle touch the bottom of the nose.


Step 4: Press the plunger



[Student Name, DOB, Student ID]

- Press the plunger firmly, using 1 motion, to deliver the dose of NAYZILAM. The person does not need to breathe deeply when NAYZILAM is given.
- Remove the nasal spray device from the nose after giving NAYZILAM.

If a second prescribed dose of NAYZILAM is needed because the seizure cluster is continuing 10 minutes after the first dose, give the second dose using a new nasal spray device in the other nostril, following the instructions above. (Repeat Step 1-4) [EDIT FOR INDIVIDUAL ORDERS]



AFTER ADMINISTERING NAYZILAM

- 1. Record and monitor**
 - Note the time NAYZILAM was administered.
 - Remain with the student and closely observe breathing, responsiveness, and seizure activity
- 2. Call 911 (EMS) if any of the following occur [CALLING FOR EMS AFTER MEDICATION ADMINISTRATION IS NOT A UNIVERSAL REQUIREMENT. EDIT FOR INDIVIDUAL ORDERS AND YOUR SETTING'S STANDARD OF PRACTICE.]**
 - Seizure does not stop or does not respond to rescue medication
 - Repeated seizures occur without full recovery between them
 - Breathing is difficult, abnormal, or the student does not regain consciousness
 - Seizure activity is significantly different from the student's typical pattern
 - A serious injury occurs
 - You are concerned about the student's condition for any reason
- 2. Notify**
 - parent/ guardian
 - school nurse
- 3. Post-seizure disposition [SENDING A STUDENT HOME AFTER MEDICATION ADMINISTRATION IS NOT A UNIVERSAL REQUIREMENT. EDIT FOR INDIVIDUAL ORDERS AND YOUR SETTING'S STANDARD OF PRACTICE.]**
 - *The decision for return to class or dismissal is individualized, based on:*
 - *the student's recovery,*
 - *provider orders*
 - *and nurse/parent/guardian input.*
- 4. Clean up and documentation**
 - Dispose of the used nasal spray device(s) and packaging.



[Student Name, DOB, Student ID]

- Complete all required documentation, including time of medication administration, response, and notifications.

ADDITIONAL INFORMATION

- Do not use NAYZILAM to treat more than 1 seizure cluster episode every 3 days or more than 5 episodes per month

Epilepsy Foundation. (2024). *Epilepsy.com*. <https://www.epilepsy.com/>

National Association of School Nurses, *School Nursing Evidence-Based Clinical Practice Guideline: Students with Seizures and Epilepsy—CPG Implementation Toolkit* (2023), 5.

UCB, Inc. (2023). *Healthcare professionals*. NAYZILAM®. <https://www.nayzilam.com/healthcare-professionals>