

You are cordially invited to attend a non-CME educational program
A 2L+ Combination Therapy for Adults With RRMM After a PI and an Immunomodulatory Agent

Presented by

**Gina Fries, PA
PA-C**

Strong Memorial Hospital, Rochester, NY

Speaker is a paid consultant of Johnson & Johnson

Date

**Wednesday, September 9, 2026
at 6:00 PM Eastern Time**

Location

**Jazzboline
5010 Main Street,
Amherst, New York
14226
(716) 839-2220**

**We look forward to your participation in this informative discussion. Reserve your spot now by contacting your Johnson & Johnson Representative, Stephen Taylor at
staylorj@its.jnj.com (585) 455-5044**

In adherence with PhRMA guidelines, spouses or other guests are not permitted to attend company-sponsored programs.

For all attendees, please be advised that information such as your name and the value and purpose of any educational item, meal, or other items of value you receive may be publicly disclosed. If you are licensed in any state or other jurisdiction, or are an employee or contractor of any organization or governmental entity that limits or prohibits meals from pharmaceutical companies, please identify yourself so that you (and we) are able to comply with such requirements.

Please note that the company prohibits the offering of gifts, gratuities, or meals to federal government employees/officials. Thank you for your cooperation.

This promotional educational activity is not accredited. The program content is developed by Johnson & Johnson. Speakers are paid consultants presenting on behalf of the company and are required to present information in compliance with FDA requirements for communications about its medicines.

The personal information you provide may be used to contact you about your request to attend Johnson & Johnson's educational program using your preferred method of communication as indicated by you. This information may be shared with Johnson & Johnson, its affiliates, and a third party for the sole purpose of completing your registration for this program and as required by law.

INDICATIONS AND USAGE

TECVAYLI[®] (teclistamab-cqyv) is a bispecific B-cell maturation antigen (BCMA)-directed CD3 T-cell engager indicated for the treatment of adult patients with relapsed or refractory multiple myeloma:

- in combination with daratumumab and hyaluronidase-fihj in patients who have received at least one prior line of therapy, including a proteasome inhibitor and an immunomodulatory agent.
- as monotherapy, in patients who have received at least four prior lines of therapy, including a proteasome inhibitor, an immunomodulatory agent and an anti-CD38 monoclonal antibody.

IMPORTANT SAFETY INFORMATION

WARNING: CYTOKINE RELEASE SYNDROME and NEUROLOGIC TOXICITY including IMMUNE EFFECTOR CELL-ASSOCIATED NEUROTOXICITY SYNDROME

Cytokine release syndrome (CRS), including life-threatening or fatal reactions, can occur in patients receiving TECVAYLI[®]. Initiate treatment with TECVAYLI[®] step-up dosing schedule to reduce risk of CRS. Withhold TECVAYLI[®] until CRS resolves or permanently discontinue based on severity.

Neurologic toxicity, including Immune Effector Cell-Associated Neurotoxicity Syndrome (ICANS) and serious, life-threatening or fatal reactions, can occur in patients receiving TECVAYLI[®]. Monitor patients for signs or symptoms of neurologic toxicity, including ICANS, during treatment. Withhold TECVAYLI[®] until neurologic toxicity resolves or permanently discontinue based on severity.

TECVAYLI[®] is available only through a restricted program called the TECVAYLI[®] and TALVEY[®] Risk Evaluation and Mitigation Strategy (REMS).

Please see the following pages for additional Important Safety Information.

Please read full [Prescribing Information](#), including Boxed WARNING, for TECVAYLI[®].

Johnson & Johnson

© Johnson & Johnson and its affiliates 2025 12/25 cp-545906v1

IMPORTANT SAFETY INFORMATION (cont.)

WARNINGS AND PRECAUTIONS

Cytokine Release Syndrome – TECVAYLI® can cause cytokine release syndrome (CRS), including life-threatening or fatal reactions.

In the clinical trials (monotherapy and combination therapy; N=448), CRS occurred in 64% of patients who received TECVAYLI® at the recommended dosage, with Grade 1 CRS occurring in 46% of patients, Grade 2 in 18%, and Grade 3 in 0.2%. Recurrent CRS occurred in 27% of patients. Most patients experienced CRS during the initial step-up dosing schedule (step-up dose 1 [37%], step-up dose 2 [32%], or the initial treatment dose [20%]). CRS first occurred following subsequent doses of TECVAYLI® in 2.5% of patients. The median time to onset of CRS was 2 (range: 1 to 9) days after the most recent dose and the median duration of CRS was 2 (range: 1 to 22) days.

Clinical signs and symptoms of CRS included, but were not limited to, fever, hypoxia, chills, hypotension, sinus tachycardia, headache, and elevated liver enzymes (aspartate aminotransferase and alanine aminotransferase elevation).

Initiate therapy according to TECVAYLI® step-up dosing schedule to reduce risk of CRS. Administer pretreatment medications to reduce risk of CRS and monitor patients following administration of TECVAYLI® accordingly.

At the first sign of CRS, immediately evaluate the patient for hospitalization. Administer supportive care based on severity and consider further management per current practice guidelines. Withhold until CRS resolves or permanently discontinue TECVAYLI® based on severity.

TECVAYLI® is available only through a restricted program under a REMS.

Neurologic Toxicity including Immune Effector Cell-Associated Neurotoxicity Syndrome - TECVAYLI® can cause serious, life-threatening, or fatal neurologic toxicity, including immune effector cell-associated neurotoxicity syndrome (ICANS).

In the clinical trials (monotherapy and combination therapy trials; N=448), neurologic toxicity occurred in 60% of patients who received TECVAYLI® at the recommended dosage, with Grade 3 or 4 neurologic toxicity in 6%. Neurologic toxicities reported in ≥5% of patients included headache (27%), sensory neuropathy (16%), motor dysfunction (15%), insomnia (12%), encephalopathy (11%), and dizziness (8%). Fatal neurologic toxicity occurred in 0.4% of patients, including Guillain-Barré syndrome and status epilepticus (one patient each).

In MajesTEC-1, ICANS was reported in 6% of patients who received TECVAYLI® as monotherapy at the recommended dosage. Recurrent ICANS occurred in 1.8% of patients. Most patients experienced ICANS following step-up dose 1 (1.2%), step-up dose 2 (0.6%), or the initial treatment dose (1.8%). Less than 3% of patients developed first occurrence of ICANS following subsequent TECVAYLI® doses. The median time to onset of ICANS was 4 (range: 2 to 8) days after the most recent dose with a median duration of 3 (range: 1 to 20) days. The most frequent clinical manifestations of ICANS reported were confusional state and dysgraphia.

In MajesTEC-3, ICANS was reported in 1.1% of patients who received the recommended TECVAYLI® dosage in combination with daratumumab and hyaluronidase-fihj, including Grade 4 ICANS in 1 patient. All events of ICANS occurred during the step-up dosing schedule. The median time to onset of ICANS was 2 (range: 1 to 3) days after the most recent dose and the median duration of ICANS was 2 (range: 1 to 2) days. The clinical manifestations of ICANS reported were amnesia, encephalopathy and delirium.

The onset of ICANS can be concurrent with CRS, following resolution of CRS, or in the absence of CRS.

Monitor patients for signs and symptoms of neurologic toxicity, including ICANS during TECVAYLI® treatment. At the first sign of neurologic toxicity, including ICANS, immediately evaluate patient and provide supportive therapy based on severity. Withhold until neurologic toxicity resolves or permanently discontinue TECVAYLI® based on severity per recommendations and consider further management per current practice guidelines.

Due to the potential for neurologic toxicity, patients receiving TECVAYLI are at risk of depressed level of consciousness. Advise patients to refrain from driving or operating heavy or potentially dangerous machinery during and for 48 hours after completion of TECVAYLI® step-up dosing schedule and in the event of new onset of any neurologic toxicity symptoms until neurologic toxicity resolves.

TECVAYLI® is available only through a restricted program under a REMS.

TECVAYLI® and TALVEY® REMS - TECVAYLI is available only through a restricted program under a REMS called the TECVAYLI® and TALVEY® REMS because of the risks of CRS and neurologic toxicity, including ICANS.

Hepatotoxicity - TECVAYLI® can cause hepatotoxicity, including fatalities. There was one fatal case of hepatic failure in MajesTEC-1. In patients who received TECVAYLI® at the recommended dosage in the clinical trials (monotherapy and combination therapy trials; N=448) elevated aspartate aminotransferase (AST) occurred in 47% of patients, with Grade 3 or 4 elevations in 2.9%. Elevated alanine aminotransferase (ALT) occurred in 48% of patients, with Grade 3 or 4 elevations in 3.8%. Elevated total bilirubin occurred in 10% of patients with Grade 3 or 4 elevations in 0.7%. Liver enzyme elevation can occur with or without concurrent CRS.

Monitor liver enzymes and bilirubin at baseline and during treatment as clinically indicated. Withhold TECVAYLI® or consider permanent discontinuation of TECVAYLI® based on severity.

Infections - TECVAYLI® can cause severe, life-threatening, or fatal infections.

In MajesTEC-1 (N=165), in patients who received the recommended TECVAYLI® dosage, serious infections, including opportunistic infections, occurred in 30% of patients, Grade 3 or 4 infections in 35% of patients, and fatal infections in 4.2% of patients.

In MajesTEC-3 (N=283), in patients who received TECVAYLI® in combination with daratumumab and hyaluronidase-fihj at the recommended dosage, serious infections, including opportunistic infections, occurred in 54% of patients, Grade 3 or Grade 4 infections in 54% of patients, and fatal infections in 4.6% of patients.

Monitor patients for signs and symptoms of infection prior to and during treatment with TECVAYLI® and treat appropriately. Administer prophylactic antimicrobials according to current practice guidelines.

Withhold TECVAYLI® or consider permanent discontinuation of TECVAYLI® based on severity.

Monitor immunoglobulin levels prior to and during treatment with TECVAYLI® and administer subcutaneous or intravenous immunoglobulin (IVIG) to maintain the serum levels >400 mg/dL.

Please read full [Prescribing Information](#), including Boxed WARNING, for TECVAYLI®.

Johnson & Johnson

© Johnson & Johnson and its affiliates 2025 12/25 cp-545906v1

IMPORTANT SAFETY INFORMATION (cont.)

Neutropenia - TECVAYLI can cause neutropenia and febrile neutropenia. In patients who received TECVAYLI® at the recommended dosage in the clinical trials (monotherapy and combination therapy trials; N=448), decreased neutrophils occurred in 88% of patients, with Grade 3 or 4 decreased neutrophils in 70%. Febrile neutropenia occurred in 6% of patients.

Monitor complete blood cell counts at baseline and periodically during treatment and provide supportive care per local institutional guidelines.

Monitor patients with neutropenia for signs of infection.

Withhold TECVAYLI® based on severity.

Hypersensitivity and Other Administration Reactions - TECVAYLI® can cause both systemic administration-related and local injection-site reactions.

Systemic Reactions - In patients who received the recommended TECVAYLI® dosage in the clinical trials (monotherapy and combination therapy trials; N=448), 2.5% of patients experienced systemic-administration reactions, which included recurrent pyrexia and rash.

Local Reactions - In patients who received TECVAYLI® at the recommended dosage in the clinical trials (monotherapy and combination therapy trials; N=448), injection-site reactions occurred in 37% of patients, with Grade 1 injection-site reactions in 29% and Grade 2 in 9%.

Withhold TECVAYLI® or consider permanent discontinuation of TECVAYLI® based on severity.

Embryo-Fetal Toxicity - Based on its mechanism of action, TECVAYLI® may cause fetal harm when administered to a pregnant patient. Advise pregnant patients of the potential risk to the fetus. Advise females of reproductive potential to use effective contraception during treatment with TECVAYLI® and for 5 months after the last dose.

ADVERSE REACTIONS

The most common adverse reactions (≥20%) in patients who received TECVAYLI® monotherapy were pyrexia, CRS, musculoskeletal pain, injection site reaction, fatigue, upper respiratory tract infection, nausea, headache, pneumonia, and diarrhea. The most common adverse reactions (≥20%) in patients who received TECVAYLI® in combination with daratumumab and hyaluronidase-fihj were hypogammaglobulinemia, upper respiratory tract infection, CRS, cough, diarrhea, musculoskeletal pain, COVID-19, pneumonia, injection site reaction, fatigue, pyrexia, headache, nausea, gastroenteritis, and weight decreased.

The most common Grade 3 to 4 laboratory abnormalities (≥20%) with TECVAYLI® (as monotherapy or in combination with daratumumab and hyaluronidase-fihj) were decreased lymphocytes, decreased neutrophils, decreased white blood cells, decreased hemoglobin, and decreased platelets.

Please read full [Prescribing Information](#), including **Boxed WARNING**, for TECVAYLI®.

cp-322928v6

Please read full [Prescribing Information](#), including **Boxed WARNING**, for TECVAYLI®.

Johnson & Johnson

© Johnson & Johnson and its affiliates 2025 12/25 cp-545906v1