

On behalf of Jazz Pharmaceuticals
You are invited to attend an expert-led discussion

ZIIHERA (Zanidatamab-hrii) for the Treatment of Adults with Previously Treated, Unresectable or Metastatic HER2-Positive (IHC 3+) Biliary Tract Cancer

Meeting Objectives:

- Understand the mechanism of disease that drives poor prognosis in biliary tract cancer (BTC)
- Explore how ZIIHERA targets HER2 to deliver dual-targeted, non-T-cell-engaging, antitumor activity
- Discover the clinical profile for the only FDA-approved, chemotherapy-free, HER2-directed 2L treatment for adult patients with unresectable or metastatic HER2+ BTC

Indication

ZIIHERA (zanidatamab-hrii) 50 mg/mL for Injection for IV is indicated for the treatment of adults with previously treated, unresectable or metastatic HER2-positive (IHC 3+) biliary tract cancer (BTC), as detected by an FDA-approved test.

This indication is approved under accelerated approval based on overall response rate and duration of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial(s).

IMPORTANT SAFETY INFORMATION

WARNING: EMBRYO-FETAL TOXICITY

Exposure to ZIIHERA during pregnancy can cause embryo-fetal harm.
Advise patients of the risk and need for effective contraception.



Program Details:

3/23/2026
5:30 PM Eastern



Featuring Faculty Presenter:

Sameh Mikhail, MD
The Mark H. Zangmeister Center



Program Location:

The Refectory Restaurant
1092 Bethel Rd.
Columbus Ohio, 43220

To register for this event, follow the registration link (if applicable) or reach out to the program host listed below.

Bertie Fields

614-581-2177 | bertie.fields@jazzpharma.com

IMPORTANT SAFETY INFORMATION (cont.)

WARNING AND PRECAUTIONS

Embryo-Fetal Toxicity

ZIIHERA can cause fetal harm when administered to a pregnant woman. In literature reports, use of a HER2-directed antibody during pregnancy resulted in cases of oligohydramnios and oligohydramnios sequence manifesting as pulmonary hypoplasia, skeletal abnormalities, and neonatal death.

Verify the pregnancy status of females of reproductive potential prior to the initiation of ZIIHERA. Advise pregnant women and females of reproductive potential that exposure to ZIIHERA during pregnancy or within 4 months prior to conception can result in fetal harm. Advise females of reproductive potential to use effective contraception during treatment with ZIIHERA and for 4 months following the last dose of ZIIHERA.

Please see reverse side for additional Important Safety Information and accompanying full Prescribing Information, including BOXED Warning.


ZIIHERA[®]
(zanidatamab-hrii)
50mg/ml Injection for IV

IMPORTANT SAFETY INFORMATION (cont.)

WARNING AND PRECAUTIONS

Left Ventricular Dysfunction

ZIIHERA can cause decreases in left ventricular ejection fraction (LVEF). LVEF declined by >10% and decreased to <50% in 4.3% of 233 patients. Left ventricular dysfunction (LVD) leading to permanent discontinuation of ZIIHERA was reported in 0.9% of patients. The median time to first occurrence of LVD was 5.6 months (range: 1.6 to 18.7). LVD resolved in 70% of patients.

Assess LVEF prior to initiation of ZIIHERA and at regular intervals during treatment. Withhold dose or permanently discontinue ZIIHERA based on severity of adverse reactions.

The safety of ZIIHERA has not been established in patients with a baseline ejection fraction that is below 50%.

Infusion-Related Reactions

ZIIHERA can cause infusion-related reactions (IRRs). An IRR was reported in 31% of 233 patients treated with ZIIHERA as a single agent in clinical studies, including Grade 3 (0.4%), and Grade 2 (25%). IRRs leading to permanent discontinuation of ZIIHERA were reported in 0.4% of patients. IRRs occurred on the first day of dosing in 28% of patients; 97% of IRRs resolved within one day.

Prior to each dose of ZIIHERA, administer premedications to prevent potential IRRs. Monitor patients for signs and symptoms of IRR during ZIIHERA administration and as clinically indicated after completion of infusion. Have medications and emergency equipment to treat IRRs available for immediate use.

If an IRR occurs, slow, or stop the infusion, and administer appropriate medical management. Monitor patients until complete resolution of signs and symptoms before resuming. Permanently discontinue ZIIHERA in patients with recurrent severe or life-threatening IRRs.

Diarrhea

ZIIHERA can cause severe diarrhea.

Diarrhea was reported in 48% of 233 patients treated in clinical studies, including Grade 3 (6%) and Grade 2 (17%). If diarrhea occurs, administer antidiarrheal treatment as clinically indicated. Perform diagnostic tests as clinically indicated to exclude other causes of diarrhea. Withhold or permanently discontinue ZIIHERA based on severity.

ADVERSE REACTIONS

Serious adverse reactions occurred in 53% of 80 patients with unresectable or metastatic HER2-positive BTC who received ZIIHERA. Serious adverse reactions in >2% of patients included biliary obstruction (15%), biliary tract infection (8%), sepsis (8%), pneumonia (5%), diarrhea (3.8%), gastric obstruction (3.8%), and fatigue (2.5%). A fatal adverse reaction of hepatic failure occurred in one patient who received ZIIHERA.

The most common adverse reactions in 80 patients with unresectable or metastatic HER2-positive BTC who received ZIIHERA ($\geq 20\%$) were diarrhea (50%), infusion-related reaction (35%), abdominal pain (29%), and fatigue (24%).

USE IN SPECIFIC POPULATIONS

Pediatric Use

Safety and efficacy of ZIIHERA have not been established in pediatric patients.

Geriatric Use

Of the 80 patients who received ZIIHERA for unresectable or metastatic HER2-positive BTC, there were 39 (49%) patients 65 years of age and older. Thirty-seven (46%) were aged 65-74 years old and 2 (3%) were aged 75 years or older.

No overall differences in safety or efficacy were observed between these patients and younger adult patients.

Please see accompanying full [Prescribing Information](#), including **BOXED warning.**

Please note that there are no certified continuing medical education credits approved for this program. Jazz Pharmaceuticals is committed to the principles of the Pharmaceutical Research and Manufacturers of America (PhRMA) Code on Interactions with Healthcare Professionals. As such, spouses and guests are not permitted to attend the program or events, and their associated expenses will not be covered or reimbursed. Due to state regulations, physicians and healthcare professionals licensed in Minnesota and Vermont are not able to receive a meal with the program. We appreciate your cooperation in this regard. Please note that meals may be reported based on various state and federal laws.

REFERENCE:

ZIIHERA (zanidatamab-hrii) Prescribing Information. Jazz Pharmaceuticals; 2024.