


YOU'RE INVITED



Please join us to learn about EMRELIS™ (telisotuzumab vedotin-tllv), the first and only c-Met targeted ADC for patients with 2L+ locally advanced or metastatic non-squamous NSCLC with high c-Met protein overexpression (3+, $\geq 50\%$).

INDICATION

EMRELIS is indicated for the treatment of adult patients with locally advanced or metastatic, non-squamous non-small cell lung cancer (NSCLC) with high c-Met protein overexpression [$\geq 50\%$ of tumor cells with strong (3+) staining], as determined by an FDA-approved test, who have received a prior systemic therapy.

This indication is approved under accelerated approval based on overall response rate (ORR) and duration of response (DOR). Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial(s).

- **Recognize** the unmet need for patients with high c-Met protein overexpressed NSCLC
- **Explore** high c-Met protein overexpression (3+, $\geq 50\%$) as an actionable biomarker in NSCLC
- **Discuss** strategies for high c-Met protein overexpression testing
- **Review** EMRELIS and identify patients who may be eligible for treatment
- **Learn** about the efficacy and safety profile of EMRELIS
- **Discuss** practical implications for EMRELIS, including dosing, dose modification, and administration



DAY AND TIME:

Tuesday, July 28, 2026
Registration: 5:30 PM CT
Presentation: 6:30 PM CT



LOCATION:

The Mercury
11909 Preston Road
Dallas, TX 75230



PLEASE RSVP BY:

Thursday, July 23, 2026
Alison Louchart
9453102522
alison.louchart@abbvie.com
<https://rsvp.abbviespeakerprograms.com/id/247836>



Keegan Russell, FNPC

Nurse Practitioner
Ironwood Cancer & Research Center
Phoenix, AZ



Scan the QR code to
register for our program

IMPORTANT SAFETY INFORMATION

Peripheral Neuropathy

EMRELIS can cause peripheral neuropathy, including peripheral sensory neuropathy and peripheral motor neuropathy. In the safety population, peripheral neuropathy occurred in 51% of patients treated with EMRELIS, including Grade 3 in 11%. These adverse reactions included peripheral sensory neuropathy in 45% of patients and peripheral motor neuropathy in 9%. The median time to onset of peripheral neuropathy was 105 days (range: 1 to 472 days). Peripheral neuropathy led to permanent discontinuation of EMRELIS in 13% of patients. The median time to onset of peripheral neuropathy leading to treatment discontinuation was 249 days (range: 57 to 519 days). Of the 7 patients with motor neuropathy ongoing as of their last dose of EMRELIS, 6 had persistent Grade 1 or 2 symptoms 30 days after their last dose.

Monitor patients for signs and symptoms of new or worsening peripheral neuropathy such as hypoesthesia, hyperesthesia, paresthesia, a burning sensation, neuropathic pain, or muscle weakness. Withhold, reduce the dose, or permanently discontinue EMRELIS based on severity.

Please see additional Important Safety Information on page 2.

Please see accompanying full Prescribing Information or visit https://www.rxabbvie.com/pdf/emrelis_pi.pdf

2L+=second-line and later. ADC=antibody-drug conjugate. c-Met=mesenchymal-epithelial transition factor. FDA=US Food and Drug Administration.

In accordance with the PhRMA Code on Interactions with Healthcare Professionals, attendance at this program is limited to healthcare professionals who practice in relevant specialties.

AbbVie tracks and reports payments and transfers of value to healthcare professionals under applicable state and federal reporting obligations.

Attendance at an AbbVie virtual webinar does not include a meal, and no other resources will be sent to you as a result of attending.

AbbVie does not report any transfer of value for attendees of its virtual webinar programs where no meal or other transfer of value is provided.



IMPORTANT SAFETY INFORMATION (CONT.)

Interstitial Lung Disease/Pneumonitis

EMRELIS can cause severe, life-threatening, or fatal interstitial lung disease (ILD)/pneumonitis. In the safety population, ILD/pneumonitis occurred in 10% of patients treated with EMRELIS, including Grade 3 in 3% and Grade 4 in 0.6%. There were 3 fatal cases of ILD/pneumonitis in patients who received EMRELIS. The median time to onset of ILD/pneumonitis was 48 days (range: 23 to 85 days). ILD/pneumonitis led to permanent discontinuation of EMRELIS in 7% of patients. The median time to onset of ILD/pneumonitis leading to treatment discontinuation was 46 days (range: 23 to 85 days).

Advise patients to immediately report cough, dyspnea, fever, and/or any new or worsening respiratory symptoms. Monitor patients for signs and symptoms of ILD/pneumonitis. Withhold or permanently discontinue EMRELIS based on severity.

Ocular Surface Disorders

EMRELIS can cause ocular surface disorders, including blurred vision, visual impairment, keratitis, and dry eye. In the safety population, ocular surface disorders occurred in 25% of patients treated with EMRELIS. The most common ocular surface disorders were blurred vision (15%), keratitis (11%), and dry eye (5%). Grade 3 ocular surface disorders occurred in 1.2% of patients [blurred vision (1.2%), and keratitis (0.6%)]. The median time to onset of ocular surface disorders was 47 days (range: 1 to 319 days).

Monitor patients for ocular surface disorders during treatment with EMRELIS. Withhold EMRELIS and refer patients to an eye care professional for an ophthalmic examination and treatment for patients who develop Grade ≥ 2 ocular toxicity. Withhold or permanently discontinue EMRELIS based on severity.

Infusion-Related Reactions

EMRELIS can cause infusion-related reactions (IRR); signs and symptoms of IRR include dyspnea, flushing, chills, nausea, chest discomfort, and hypotension. The median time to onset of IRR was 28 days (range: 1 to 43 days). In the safety population, IRR occurred in 3% of patients treated with EMRELIS, including Grade 3 in 1.2% and Grade 4 in 0.6%. IRR led to permanent discontinuation of EMRELIS in 0.6% of patients.

Monitor patients for signs and symptoms of infusion reactions during EMRELIS infusion. Withhold, reduce the rate of infusion, or permanently discontinue EMRELIS based on severity. For patients who experience IRR, administer premedications prior to subsequent infusions.

Embryo-Fetal Toxicity

Based on the mechanism of action and findings in animals, EMRELIS can cause fetal harm when administered to a pregnant woman. The small molecule component of EMRELIS, monomethyl auristatin E (MMAE), administered to rats caused adverse developmental outcomes, including embryo-fetal mortality and structural abnormalities, at exposures similar to those occurring clinically at the recommended dose.

Advise patients of the potential risk to a fetus. Advise females of reproductive potential to use effective contraception during treatment with EMRELIS and for 2 months after the last dose. Advise male patients with female partners of reproductive potential to use effective contraception during treatment with EMRELIS and for 4 months after the last dose.

Adverse Reactions

Serious adverse reactions occurred in 35% of patients. The most common adverse reactions ($\geq 20\%$) were peripheral neuropathy, fatigue, decreased appetite, and peripheral edema.

The most common Grade 3 or 4 laboratory abnormalities ($\geq 2\%$) were decreased lymphocytes, increased glucose, increased alanine aminotransferase, increased gamma glutamyl transferase, decreased phosphorus, decreased sodium, decreased hemoglobin, and decreased calcium.

Drug Interactions

Strong CYP3A Inhibitors: Concomitant use with EMRELIS may increase the area under the curve of MMAE. Monitor for increased risk of adverse reactions to EMRELIS.

Use in Specific Populations

Severe or Moderate Hepatic Impairment: Avoid the use of EMRELIS.

Lactation: Advise lactating women not to breastfeed during treatment with EMRELIS and for 1 month after the last dose.

Infertility: Based on findings from animal studies, EMRELIS may impair fertility in females and males.

Please see accompanying full Prescribing Information or visit https://www.rxabbvie.com/pdf/emrelis_pi.pdf



Emrelis™
(telisotuzumab vedotin-tllv)
injection for intravenous use 20 mg/100 mg