

# Please join us for a live presentation on LIBTAYO® (cemiplimab-rwlc)

## FDA APPROVED IN ADVANCED BASAL CELL CARCINOMA (BCC) AND ADVANCED CUTANEOUS SQUAMOUS CELL CARCINOMA (CSCC)

### LIBTAYO (cemiplimab-rwlc) is:

- The first and only treatment indicated for patients with locally advanced or metastatic basal cell carcinoma (laBCC or mBCC) who have been previously treated with a hedgehog pathway inhibitor (HHI) or for whom an HHI is not appropriate
- The first treatment indicated for patients with metastatic cutaneous squamous cell carcinoma (mCSCC) or locally advanced CSCC (laCSCC) who are not candidates for curative surgery or curative radiation

#### PROGRAM AGENDA

5/9/2024

6:00 PM Central Presentation

(Please arrive 30 minutes prior for registration)

#### PRESENTED BY

Grace Cherry, NP  
UCLA Medical Group  
Los Angeles, CA

#### LOCATION

Bob's Steak & Chop House  
5760 Legacy Drive  
Plano, Texas 75024

#### HOSTED BY

Teresa Hill  
teresa.hill@regeneron.com  
469-343-8810  
RGN0030962  
*You may RSVP to your program host.*

To register for this event, please call 469-343-8810  
and reference meeting ID RGN0030962, or visit <https://tppprod-regeneron.physiciansworld.com/9tO4EA>.

#### DURING THIS PRESENTATION, WE WILL DISCUSS:

- Utilizing a multidisciplinary team for managing patients with advanced BCC and advanced CSCC
- The clinical efficacy and safety data of LIBTAYO in advanced BCC and advanced CSCC
- LIBTAYO patient case studies

#### Important Safety Information

##### Warnings and Precautions

###### Severe and Fatal Immune-Mediated Adverse Reactions

Immune-mediated adverse reactions, which may be severe or fatal, can occur in any organ system or tissue at any time after starting treatment. While immune-mediated adverse reactions usually occur during treatment, they can also occur after discontinuation. Immune-mediated adverse reactions affecting more than one body system can occur simultaneously. Early identification and management are essential to ensuring safe use of PD-1/PD-L1-blocking antibodies. The definition of immune-mediated adverse reactions included the required use of systemic corticosteroids or other immunosuppressants and the absence of a clear alternate etiology. Monitor closely for symptoms and signs that may be clinical manifestations of underlying immune-mediated adverse reactions. Evaluate liver enzymes, creatinine, and thyroid function at baseline and periodically during treatment. In cases of suspected immune-mediated adverse reactions, initiate appropriate workup to exclude alternative etiologies, including infection. Institute medical management promptly, including specialty consultation as appropriate.

No dose reduction for LIBTAYO is recommended. In general, withhold LIBTAYO for severe (Grade 3) immune-mediated adverse reactions. Permanently discontinue LIBTAYO for life-threatening (Grade 4) immune-mediated adverse reactions, recurrent severe (Grade 3) immune-mediated adverse reactions that require systemic immunosuppressive treatment, or an inability to reduce corticosteroid dose to 10 mg or less of prednisone equivalent per day within 12 weeks of initiating steroids.

Withhold or permanently discontinue LIBTAYO depending on severity. In general, if LIBTAYO requires interruption or discontinuation, administer systemic corticosteroid therapy (1 to 2 mg/kg/day prednisone or equivalent) until improvement to Grade 1 or less. Upon improvement to Grade 1 or less, initiate corticosteroid taper and continue to taper over at least 1 month. Consider administration of other systemic immunosuppressants in patients whose immune-mediated adverse reactions are not controlled with corticosteroids.

The incidence and severity of immune-mediated adverse reactions were similar when LIBTAYO was administered as a single agent or in combination with chemotherapy.

Please see additional Important Safety Information throughout and accompanying full [Prescribing Information](#).

In accordance with the PhRMA Code on Interactions with Healthcare Professionals, this Program is limited to U.S. Healthcare Professionals and persons with bona fide professional interest in the information presented. Attendance at this Program by guests or spouses is not permitted unless they would qualify as an appropriate attendee on their own. If a meal is provided, actively licensed Minnesota and Vermont prescribers may attend, but not partake in the meal. Full-time Federal Employees may attend and partake in the meal if the Program is considered widely attended (50 or more attendees). If not, they may attend, but not partake in the meal. Part-time Federal Employees acting in their civilian capacity may attend the Program and partake in the meal. Alcohol is not permitted in connection with Speaker Programs and will not be provided. The value of any meal provided in connection with the Program may be reported in accordance with federal and state laws and regulations.

## Important Safety Information (cont'd)

### Warnings and Precautions (cont'd)

**Severe and Fatal Immune-Mediated Adverse Reactions (cont'd)**

**Immune-mediated pneumonitis:** LIBTAYO can cause immune-mediated pneumonitis. In patients treated with other PD-1/PD-L1-blocking antibodies, the incidence of pneumonitis is higher in patients who have received prior thoracic radiation. Immune-mediated pneumonitis occurred in 2.6% (33/1281) of patients receiving LIBTAYO, including Grade 4 (0.3%), Grade 3 (0.6%), and Grade 2 (1.6%). Pneumonitis led to permanent discontinuation in 1.3% of patients and withholding of LIBTAYO in 1.4% of patients. Systemic corticosteroids were required in all patients with pneumonitis. Pneumonitis resolved in 61% of the 33 patients. Of the 18 patients in whom LIBTAYO was withheld, 10 reinitiated after symptom improvement; of these, 4/10 (40%) had recurrence of pneumonitis. Withhold LIBTAYO for Grade 2, and permanently discontinue for Grade 3 or 4. Resume in patients with complete or partial resolution (Grade 0 to 1) after corticosteroid taper. Permanently discontinue if no complete or partial resolution within 12 weeks of initiating steroids or inability to reduce prednisone to less than 10 mg per day (or equivalent) within 12 weeks of initiating steroids.

**Immune-mediated colitis:** LIBTAYO can cause immune-mediated colitis. The primary component of immune-mediated colitis was diarrhea. Cytomegalovirus (CMV) infection/reactivation has been reported in patients with corticosteroid-refractory immune-mediated colitis treated with PD-1/PD-L1-blocking antibodies. In cases of corticosteroid-refractory immune-mediated colitis, consider repeating infectious workup to exclude alternative etiologies. Immune-mediated colitis occurred in 2% (25/1281) of patients receiving LIBTAYO, including Grade 3 (0.8%) and Grade 2 (0.9%). Colitis led to permanent discontinuation in 0.4% of patients and withholding of LIBTAYO in 1.2% of patients. Systemic corticosteroids were required in all patients with colitis. Colitis resolved in 56% of the 25 patients. Of the 16 patients in whom LIBTAYO was withheld, 6 reinitiated LIBTAYO after symptom improvement; of these, 4/6 (67%) had recurrence. Withhold LIBTAYO for Grade 2 or 3, and permanently discontinue for Grade 4. Resume in patients with complete or partial resolution (Grade 0 to 1) after corticosteroid taper. Permanently discontinue if no complete or partial resolution within 12 weeks of initiating steroids or inability to reduce prednisone to less than 10 mg per day (or equivalent) within 12 weeks of initiating steroids.

**Immune-mediated hepatitis:** LIBTAYO can cause immune-mediated hepatitis. Immune-mediated hepatitis occurred in 2.4% (31/1281) of patients receiving LIBTAYO, including fatal (<0.1%), Grade 4 (0.3%), Grade 3 (1.6%), and Grade 2 (0.2%). Hepatitis led to permanent discontinuation of LIBTAYO in 1.4% of patients and withholding of LIBTAYO in 0.7% of patients. Systemic corticosteroids were required in all patients with hepatitis. Additional immunosuppression with mycophenolate was required in 13% (4/31) of these patients. Hepatitis resolved in 39% of the 31 patients. Of the 9 patients in whom LIBTAYO was withheld, 5 reinitiated LIBTAYO after symptom improvement; of these, 1/5 (20%) had recurrence.

**For hepatitis with no tumor involvement of the liver:** Withhold LIBTAYO if AST or ALT increases to more than 3 and up to 8 times the upper limit of normal (ULN) or if total bilirubin increases to more than 1.5 and up to 3 times the ULN. Permanently discontinue LIBTAYO if AST or ALT increases to more than 8 times the ULN or total bilirubin increases to more than 3 times the ULN.

**For hepatitis with tumor involvement of the liver:** Withhold LIBTAYO if baseline AST or ALT is more than 1 and up to 3 times ULN and increases to more than 5 and up to 10 times ULN. Also, withhold LIBTAYO if baseline AST or ALT is more than 3 and up to 5 times ULN and increases to more than 8 and up to 10 times ULN. Permanently discontinue LIBTAYO if AST or ALT increases to more than 10 times ULN or if total bilirubin increases to more than 3 times ULN. If AST and ALT are less than or equal to ULN at baseline, withhold or permanently discontinue LIBTAYO based on recommendations for hepatitis with no liver involvement.

Resume in patients with complete or partial resolution (Grade 0 to 1) after corticosteroid taper. Permanently discontinue if no complete or partial resolution within 12 weeks of initiating steroids or inability to reduce prednisone to less than 10 mg per day (or equivalent) within 12 weeks of initiating steroids.

**Immune-mediated endocrinopathies:** For Grade 3 or 4 endocrinopathies, withhold until clinically stable or permanently discontinue depending on severity.

**• Adrenal insufficiency:** LIBTAYO can cause primary or secondary adrenal insufficiency. For Grade 2 or higher adrenal insufficiency, initiate symptomatic treatment, including hormone replacement as clinically indicated. Withhold LIBTAYO depending on severity. Adrenal insufficiency occurred in 0.5% (6/1281) of patients receiving LIBTAYO, including Grade 3 (0.5%). Adrenal insufficiency led to permanent discontinuation of LIBTAYO in 1 (<0.1%) patient. LIBTAYO was withheld in 1 (<1%) patient due to adrenal insufficiency and not reinitiated. Systemic corticosteroids were required in 83% (5/6) patients with adrenal insufficiency; of these, the majority remained on systemic corticosteroids. Adrenal insufficiency had resolved in 17 of the 6 patients

**• Hypophysitis:** LIBTAYO can cause immune-mediated hypophysitis. Hypophysitis can present with acute symptoms associated with mass effect such as headache, photophobia, or visual field defects. Hypophysitis can cause hypopituitarism. Initiate hormone replacement as clinically indicated. Withhold or permanently discontinue depending on severity. Hypophysitis occurred in 0.5% (7/1281) of patients receiving LIBTAYO, including Grade 3 (0.2%) and Grade 2 (0.3%) adverse reactions. Hypophysitis led to permanent discontinuation of LIBTAYO in 1 (<0.1%) patient and withholding of LIBTAYO in 2 (0.2%) patients. Systemic corticosteroids were required in 86% (6/7) of patients with hypophysitis. Hypophysitis resolved in 14% of the 7 patients. Of the 2 patients in whom LIBTAYO was withheld for hypophysitis, none of the patients reinitiated

**• Thyroid disorders:** LIBTAYO can cause immune-mediated thyroid disorders. Thyroiditis can present with or without endocrinopathy. Hypothyroidism can follow hyperthyroidism. Initiate hormone replacement or medical management of hyperthyroidism as clinically indicated. Withhold or permanently discontinue LIBTAYO depending on severity

**• Thyroiditis:** Thyroiditis occurred in 0.6% (8/1281) of patients receiving LIBTAYO, including Grade 2 (0.3%) adverse reactions. No patient discontinued LIBTAYO due to thyroiditis. Thyroiditis led to withholding of LIBTAYO in 1 (<0.1%) patient. Systemic corticosteroids were not required in any patient with thyroiditis. Thyroiditis resolved in 13% of the 8 patients. Blood thyroid stimulating hormone increased and blood thyroid stimulating hormone decreased have also been reported

**• Hyperthyroidism:** Hyperthyroidism occurred in 3% (39/1281) of patients receiving LIBTAYO, including Grade 3 (<0.1%) and Grade 2 (0.9%). No patient discontinued treatment and LIBTAYO was withheld in 7 (0.5%) patients due to hyperthyroidism. Systemic corticosteroids were required in 8% (3/39) of patients. Hyperthyroidism resolved in 56% of 39 patients. Of the 7 patients in whom LIBTAYO was withheld for hyperthyroidism, 2 patients reinitiated LIBTAYO after symptom improvement; of these, none had recurrence of hyperthyroidism

**• Hypothyroidism:** Hypothyroidism occurred in 7% (87/1281) of patients receiving LIBTAYO, including Grade 3 (<0.1%) and Grade 2 (6%). Hypothyroidism led to permanent discontinuation of LIBTAYO in 3 (0.2%) patients. Hypothyroidism led to withholding of LIBTAYO in 9 (0.7%) patients. Systemic corticosteroids were required in 1.1% (1/87) of patients with hypothyroidism. Hypothyroidism resolved in 6% of the 87 patients. Majority of the patients with hypothyroidism required long-term thyroid hormone replacement. Of the 9 patients in whom LIBTAYO was withheld for hypothyroidism, 1 reinitiated LIBTAYO after symptom improvement and did not have recurrence of hypothyroidism

**• Type 1 diabetes mellitus, which can present with diabetic ketoacidosis:** Monitor for hyperglycemia or other signs and symptoms of diabetes. Initiate treatment with insulin as clinically indicated. Withhold LIBTAYO depending on severity. Type 1 diabetes mellitus occurred in <0.1% (1/1281) of patients (Grade 4).

Please see additional Important Safety Information throughout and accompanying full [Prescribing Information](#).

In accordance with the PhRMA Code on Interactions with Healthcare Professionals, this Program is limited to U.S. Healthcare Professionals and persons with bona fide professional interest in the information presented. Attendance at this Program by guests or spouses is not permitted unless they would qualify as an appropriate attendee on their own. If a meal is provided, actively licensed Minnesota and Vermont prescribers may attend, but not partake in the meal. Full-time Federal Employees may attend and partake in the meal if the Program is considered widely attended (50 or more attendees). If not, they may attend, but not partake in the meal. Part-time Federal Employees acting in their civilian capacity may attend the Program and partake in the meal. Alcohol is not permitted in connection with Speaker Programs and will not be provided. The value of any meal provided in connection with the Program may be reported in accordance with federal and state laws and regulations.

## Important Safety Information (cont'd)

### Warnings and Precautions (cont'd)

#### Severe and Fatal Immune-Mediated Adverse Reactions (cont'd)

##### Type 1 diabetes mellitus, which can present with diabetic ketoacidosis: (cont'd)

No patient discontinued treatment due to Type 1 diabetes mellitus. Type 1 diabetes mellitus led to withholding of LIBTAYO in 0.1% of patients, treatment was reinitiated after symptom improvement. Patient received long-term insulin therapy.

**Immune-mediated nephritis with renal dysfunction:** LIBTAYO can cause immune-mediated nephritis. Immune-mediated nephritis occurred in 0.7% (9/1281) of patients receiving LIBTAYO, including fatal (<0.1%), Grade 3 (<0.1%), and Grade 2 (0.5%). Nephritis led to permanent discontinuation in 0.2% of patients and withholding of LIBTAYO in 0.4% of patients. Systemic corticosteroids were required in all patients with nephritis. Nephritis resolved in 78% of the 9 patients. Of the 5 patients in whom LIBTAYO was withheld, 4 reinitiated LIBTAYO after symptom improvement; of these, 1/4 (25%) had recurrence. Withhold LIBTAYO for Grade 2 or 3 increased blood creatinine, and permanently discontinue for Grade 4 increased blood creatinine. Resume in patients with complete or partial resolution (Grade 0 to 1) after corticosteroid taper. Permanently discontinue if no complete or partial resolution within 12 weeks of initiating steroids or inability to reduce prednisone to less than 10 mg per day (or equivalent) within 12 weeks of initiating steroids.

**Immune-mediated dermatologic adverse reactions:** LIBTAYO can cause immune-mediated rash or dermatitis. Exfoliative dermatitis, including Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), and drug rash with eosinophilia and systemic symptoms (DRESS) has occurred with PD-1/PD-L1-blocking antibodies. Immune-mediated dermatologic adverse reactions occurred in 1.9% (24/1281) of patients receiving LIBTAYO, including Grade 3 (0.9%) and Grade 2 (0.8%).

Immune-mediated dermatologic adverse reactions led to permanent discontinuation in 0.2% of patients and withholding of LIBTAYO in 1.3% of patients. Systemic corticosteroids were required in all patients with immune-mediated dermatologic adverse reactions. Immune-mediated dermatologic adverse reactions resolved in 71% of the 24 patients. Of the 17 patients in whom LIBTAYO was withheld for dermatologic adverse reaction, 13 reinitiated LIBTAYO after symptom improvement; of these, 5/13 (38%) had recurrence of the dermatologic adverse reaction. Topical emollients and/or topical corticosteroids may be adequate to treat mild to moderate non-exfoliative rashes. Withhold LIBTAYO for suspected SJS, TEN, or DRESS. Permanently discontinue LIBTAYO for confirmed SJS, TEN, or DRESS. Resume in patients with complete or partial resolution (Grade 0 to 1) after corticosteroid taper. Permanently discontinue if no complete or partial resolution within 12 weeks of initiating steroids or inability to reduce prednisone to less than 10 mg per day (or equivalent) within 12 weeks of initiating steroids.

**Other immune-mediated adverse reactions:** The following clinically significant immune-mediated adverse reactions occurred at an incidence of <1% in 1281 patients who received LIBTAYO or were reported with the use of other PD-1/PD-L1-blocking antibodies. Severe or fatal cases have been reported for some of these adverse reactions.

- Cardiac/vascular:** Myocarditis, pericarditis, and vasculitis. Permanently discontinue for Grades 2, 3, or 4 myocarditis
- Nervous system:** Meningitis, encephalitis, myelitis and demyelination, myasthenic syndrome/myasthenia gravis (including exacerbation), Guillain-Barré syndrome, nerve paresis, and autoimmune neuropathy. Withhold for Grade 2 neurological toxicities and permanently discontinue for Grades 3 or 4 neurological toxicities. Resume in patients with complete or partial resolution (Grade 0 to 1) after corticosteroid taper. Permanently discontinue if no complete or partial resolution within 12 weeks of initiating steroids or inability to reduce prednisone to less than 10 mg per day (or equivalent) within 12 weeks of initiating steroids
- Ocular:** Uveitis, iritis, and other ocular inflammatory toxicities. Some cases can be associated with retinal detachment. Various grades of

visual impairment to include blindness can occur. If uveitis occurs in combination with other immune-mediated adverse reactions, consider a Vogt-Koyanagi-Harada-like syndrome, as this may require treatment with systemic steroids to reduce the risk of permanent vision loss.

- Gastrointestinal:** Pancreatitis to include increases in serum amylase and lipase levels, gastritis, duodenitis, stomatitis

- Musculoskeletal and connective tissue:** Myositis/polymyositis, rhabdomyolysis, and associated sequelae including renal failure, arthritis, polymyalgia rheumatica

- Endocrine:** Hypoparathyroidism

- Other (hematologic/immune):** Hemolytic anemia, aplastic anemia, hemophagocytic lymphohistiocytosis, systemic inflammatory response syndrome, histiocytic necrotizing lymphadenitis (Kikuchi lymphadenitis), sarcoidosis, immune thrombocytopenia, solid organ transplant rejection

#### Infusion-Related Reactions

Severe or life-threatening infusion-related reactions occurred in 0.2% of patients receiving LIBTAYO as a single agent. Monitor patients for signs and symptoms of infusion-related reactions. Common symptoms of infusion-related reaction include nausea, pyrexia, and vomiting. Interrupt or slow the rate of infusion or permanently discontinue LIBTAYO based on severity of reaction.

#### Complications of Allogeneic HSCT

Fatal and other serious complications can occur in patients who receive allogeneic hematopoietic stem cell transplantation (HSCT) before or after being treated with a PD-1/PD-L1-blocking antibody. Transplant-related complications include hyperacute graft-versus-host disease (GVHD), acute GVHD, chronic GVHD, hepatic veno-occlusive disease (VOD) after reduced intensity conditioning, and steroid-requiring febrile syndrome (without an identified infectious cause). These complications may occur despite intervening therapy between PD-1/PD-L1 blockade and allogeneic HSCT. Follow patients closely for evidence of transplant-related complications and intervene promptly. Consider the benefit versus risks of treatment with a PD-1/PD-L1-blocking antibody prior to or after an allogeneic HSCT.

#### Embryo-Fetal Toxicity

LIBTAYO can cause fetal harm when administered to a pregnant woman due to an increased risk of immune-mediated rejection of the developing fetus resulting in fetal death. Advise women of the potential risk to a fetus. Advise females of reproductive potential to use effective contraception during treatment with LIBTAYO and for at least 4 months after the last dose.

#### Adverse Reactions

**LIBTAYO as a single agent:** the most common adverse reactions (≥15%) are fatigue, musculoskeletal pain, rash, diarrhea, and anemia

**LIBTAYO in combination with platinum-based chemotherapy:** the most common adverse reactions (≥15%) are alopecia, musculoskeletal pain, nausea, fatigue, peripheral neuropathy, and decreased appetite

#### Use in Specific Populations

- Lactation:** Because of the potential for serious adverse reactions in breastfed children, advise women not to breastfeed during treatment and for at least 4 months after the last dose of LIBTAYO

- Females and males of reproductive potential:** Verify pregnancy status in females of reproductive potential prior to initiating LIBTAYO

Please see accompanying full [Prescriber Information](#)